

## REFERENCE

- 1 **AVID Investigators.** A comparison of antiarrhythmic-drug therapy with implantable defibrillators in patients resuscitated from near-fatal ventricular arrhythmias. The Antiarrhythmics versus implantable defibrillators (AVID) investigators. *N Engl J Med* 1997;**337**:1576–83.
- 2 **Kadish A, Dyer A, Daubert JP, et al.** Prophylactic defibrillator implantation in patients with nonischemic dilated cardiomyopathy. *N Engl J Med* 2004;**350**:2151–8.
- 3 **Moss AJ, Zareba W, Hall WJ, et al.** Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med* 2002;**346**:877–83.
- 4 **Anvari A, Gottsauner-Wolf M, Turel Z, et al.** Predictors of outcome in patients with implantable cardioverter defibrillators. *Cardiology* 1998;**90**:180–6.
- 5 **Dolack GL.** Clinical predictors of implantable cardioverter-defibrillator shocks (results of the CASCADE trial). Cardiac arrest in Seattle, conventional versus amiodarone drug evaluation. *Am J Cardiol* 1994;**73**:237–41.
- 6 **Grimm W, Flores BT, Marchlinski FE.** Shock occurrence and survival in 241 patients with implantable cardioverter-defibrillator therapy. *Circulation* 1993;**87**:1880–8.
- 7 **Levine JH, Mellits ED, Baumgardner RA, et al.** Predictors of first discharge and subsequent survival in patients with automatic implantable cardioverter-defibrillators. *Circulation* 1991;**84**:558–66.
- 8 **Rankovic V, Karha J, Passman R, et al.** Predictors of appropriate implantable cardioverter-defibrillator therapy in patients with idiopathic dilated cardiomyopathy. *Am J Cardiol* 2002;**89**:1072–6.
- 9 **Whang W, Mittleman MA, Rich DQ, et al.** Heart failure and the risk of shocks in patients with implantable cardioverter defibrillators: results from the triggers of ventricular arrhythmias (TOVA) study. *Circulation* 2004;**109**:1386–91.
- 10 **Reiter MJ, Fain ES, Senelly KM, et al.** Predictors of device activation for ventricular arrhythmias and survival in patients with implantable pacemakers/defibrillators. CADENCE investigators. *Pacing Clin Electrophysiol* 1994;**17**:1487–98.
- 11 **Freedberg NA, Hill JN, Fogel RI, et al.** Recurrence of symptomatic ventricular arrhythmias in patients with implantable cardioverter defibrillator after the first device therapy: implications for antiarrhythmic therapy and driving restrictions. CARE Group. *J Am Coll Cardiol* 2001;**37**:1910–5.
- 12 **Yin WH, Chen JW, Jen HL, et al.** Independent prognostic value of elevated high-sensitivity C-reactive protein in chronic heart failure. *Am Heart J* 2004;**147**:931–8.
- 13 **Yu CM, Sanderson JE.** Plasma brain natriuretic peptide—an independent predictor of cardiovascular mortality in acute heart failure. *Eur J Heart Fail* 1999;**1**:59–65.
- 14 **Wang TJ, Larson MG, Levy D, et al.** Plasma natriuretic peptide levels and the risk of cardiovascular events and death. *N Engl J Med* 2004;**350**:655–63.
- 15 **Tapainen JM, Lindgren KS, Makikallio TH, et al.** Natriuretic peptides as predictors of non-sudden and sudden cardiac death after acute myocardial infarction in the beta-blocking era. *J Am Coll Cardiol* 2004;**43**:757–63.
- 16 **Berger R, Huelsman M, Strecker K, et al.** B-type natriuretic peptide predicts sudden death in patients with chronic heart failure. *Circulation* 2002;**105**:2392–7.
- 17 **Isnard R, Pousset F, Chafirovskaja O, et al.** Combination of B-type natriuretic peptide and peak oxygen consumption improves risk stratification in outpatients with chronic heart failure. *Am Heart J* 2003;**146**:729–35.
- 18 **Wallen T, Landahl S, Hedner T, et al.** Brain natriuretic peptide predicts mortality in the elderly. *Heart* 1997;**77**:264–7.
- 19 **Gibelin P.** An evaluation of symptom classification systems used for the assessment of patients with heart failure in France. *Eur J Heart Fail* 2001;**3**:739–46.
- 20 **Vanderheyden M, Bartunek J, Goethals M.** Brain and other natriuretic peptides: molecular aspects. *Eur J Heart Fail* 2004;**6**:261–8.
- 21 **Hansen DE, Craig CS, Hondeghem LM.** Stretch-induced arrhythmias in the isolated canine ventricle: evidence for the importance of mechano-electrical feedback. *Circulation* 1990;**81**:1094–105.
- 22 **Zhu WX, Johnson SB, Brandt R, et al.** Impact of volume loading and load reduction on ventricular refractoriness and conduction properties in canine congestive heart failure. *J Am Coll Cardiol* 1997;**30**:825–33.
- 23 **Dogra V, Oliver R, Lapidus J, et al.** Apparent protective effect of increased left ventricular wall thickness in an ICD population. *J Card Fail* 2003;**9**:412–5.
- 24 **Albert CM, Ma J, Rifai N, et al.** Prospective study of C-reactive protein, homocysteine, and plasma lipid levels as predictors of sudden cardiac death. *Circulation* 2002;**105**:2595–9.
- 25 **Bailey JJ, Berson AS, Handelsman H, et al.** Utility of current risk stratification tests for predicting major arrhythmic events after myocardial infarction. *J Am Coll Cardiol* 2001;**38**:1902–11.
- 26 **Klein RC, Raitt MH, Wilkoff BL, et al.** Analysis of implantable cardioverter defibrillator therapy in the antiarrhythmics versus implantable defibrillators (AVID) trial. *J Cardiovasc Electrophysiol* 2003;**14**:940–8.
- 27 **Bardy GH, Lee KL, Mark DB, et al.** Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med* 2005;**352**:225–37.

## IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.068163

## Myocardial calcification: a rare cause of diastolic dysfunction

**A** 65 year old man who suffered a large anteroapical myocardial infarction 12 years previously was admitted to the cardiology ward for investigation of heart failure. A roentgen chest x ray showed calcification projecting onto the apex of the heart which corresponded on computer chest x ray to an extensive circumferential area of transmyocardial calcification of the old infarcted area (panels A (transverse view) and B (sagittal view)). Serum calcium and parathyroid hormone concentrations were normal. To assess any functional cardiac changes due to these calcifications, a trans-thoracic echocardiogram was performed, which documented a left ventricular ejection fraction of 30% (Simpson method), an akinetic apical and mid part of the left ventricle, and a persisting restrictive inflow pattern even after high dose treatment with intravenous diuretic therapy. We postulate that part of the restrictive inflow pattern is secondary due to the stiff characteristic of the ventricle mainly caused by a large area of myocardial calcification. Cardiac calcifications have been reported resulting from secondary hyperparathyroidism, most frequently related to chronic renal failure and uraemia and to atherosclerosis mainly in the



coronary arteries or cardiac valves. However, a detailed review of the literature revealed no reports of extensive circumferential myocardial calcification secondary to an old myocardial infarction.

W Mullens  
J D Keyser  
W Droogne  
wmulle0@hotmail.com